

**Amendments to the Claims**

This listing of claims replaces all prior versions, and listings, of claims in the above-identified application:

**Listing of Claims**

1. **(Currently Amended)** A method of inducing a cytokine in a cell, the method comprising contacting the cell with an immunological regulator under conditions effective to induce a cytokine, wherein the immunological regulator ~~comprises~~ consists of MQPPPLP (SEQ ID NO:1), an active analog thereof, and combinations thereof, wherein the active analog comprises a peptide having an amino acid sequence with at least about 15 percent proline and having at least about 70 percent ~~structural similarity~~ sequence identity to SEQ ID NO:1.
2. **(Original)** The method of claim 1 wherein the cell is present in a cell culture, a tissue, an organ, or an organism.
3. **(Original)** The method of claim 1 wherein the cell is a mammalian cell.
4. **(Original)** The method of claim 3 wherein the cell is a human cell.
5. **(Cancelled)**
6. **(Currently Amended)** A method for modulating an immune response in a cell, the method comprising contacting the cell with an immunological regulator under conditions effective to induce a cytokine, wherein the immunological regulator ~~comprises~~ consists of MQPPPLP (SEQ ID NO:1), an active analog thereof, and combinations thereof, wherein the active analog comprises a peptide having an amino acid sequence with at least about 15 percent

proline and having at least about 70 percent ~~structural similarity~~ sequence identity to SEQ ID NO:1.

7. **(Original)** The method of claim 6 wherein the cell is present in a cell culture, a tissue, an organ, or an organism.

8. **(Original)** The method of claim 6 wherein the cell is a mammalian cell.

9. **(Original)** The method of claim 8 wherein the cell is a human cell.

10. **(Cancelled)**

11. **(Currently Amended)** A method for modulating an immune response in a patient, the method comprising administering to the patient an immunological regulator under conditions effective to induce a cytokine, wherein the immunological regulator ~~comprises~~ consists of MQPPPLP (SEQ ID NO:1), an active analog thereof, and combinations thereof, wherein the active analog comprises a peptide having an amino acid sequence with at least about 15 percent proline and having at least about 70 percent ~~structural similarity~~ sequence identity to SEQ ID NO:1.

12. **(Cancelled)**

13. **(Original)** The method of claim 11 wherein the immunological regulator is administered as part of a dietary supplement.

14. **(Original)** The method of claim 11 wherein the immunological regulator is administered topically.

15. **(Original)** The method of claim 11 wherein the patient is an animal.
16. **(Original)** The method of claim 15 wherein the patient is a human.
17. **(Original)** The method of claim 11 wherein the immune response is a specific immune response.
18. **(Original)** The method of claim 11 wherein the immune response is a nonspecific immune response.
19. **(Original)** The method of claim 11 wherein the immune response is the interferon response or antibody production.
20. **(Currently Amended)** A method for modulating leukocyte proliferation, the method comprising contacting leukocytes with a leukocyte regulator selected from the group of colostrinin, a constituent peptide thereof, an active analog thereof, and combinations thereof, under conditions effective to change the number of leukocytes; wherein the active analog comprises a peptide having an amino acid sequence with at least about 15 percent proline and having at least about 70 percent ~~structural similarity~~ sequence identity to one or more constituent peptides of colostrinin, which are selected from the group of SEQ ID NO:1 through SEQ ID NO:34.
21. **(Previously Presented)** The method of claim 20 wherein the leukocytes are present in a cell culture or an organism.
22. **(Previously Presented)** The method of claim 20 wherein the leukocytes are mammalian cells.

23. **(Previously Presented)** The method of claim 22 wherein the leukocytes are human cells.
24. **(Previously Presented)** The method of claim 22 wherein the leukocytes are increased in number.
25. **(Previously Presented)** The method of claim 24 wherein the leukocytes are differentiated.
26. **(Previously Presented)** The method of claim 22 wherein the leukocyte regulator is a constituent peptide of colostrinin.
27. **(Currently Amended)** The method of claim 26 wherein the leukocyte regulator is selected from the group of MQPPPLP (SEQ ID NO:1), LQTPQPLLQVMMEPQGD (SEQ ID NO:2), DQPPDVEKPDQLQPFQVQS (SEQ ID NO:3), LFFFLPVVNVLP (SEQ ID NO:4), DLEMPVLPVEPPFPV (SEQ ID NO:5), MPQNFYKLPQM (SEQ ID NO:6), VLEMKFPPPPQETVT (SEQ ID NO:7), LKPFPKLKVEVFPPF (SEQ ID NO:8), VVMEV (SEQ ID NO:9), SEQP (SEQ ID NO:10), DKE (SEQ ID NO:11), FPPPK (SEQ ID NO:12), DSQPPV (SEQ ID NO:13), DPPPPQS (SEQ ID NO:14), SEEMP (SEQ ID NO:15), KYKLQPE (SEQ ID NO:16), VLPPNVG (SEQ ID NO:17), VYPFTGPIPN (SEQ ID NO:18), SLPQNILPL (SEQ ID NO:19), TQTPVVVPPF (SEQ ID NO:20), LQPEIMGVPKVKETMVPK (SEQ ID NO:21), HKEMPFPKYPVEPFTESQ (SEQ ID NO:22), SLTLTDVEKLHLPLPLVQ (SEQ ID NO:23), SWMHQPP (SEQ ID NO:24), QPLPPTVMFP (SEQ ID NO:25), PQSVLS (SEQ ID NO:26), LSQPKVLPVPQKAVPQRDMPIQ (SEQ ID NO:27), AFLLYQE (SEQ ID NO:28), RGPFPILV (SEQ ID NO:29), ATFNRYQDDHGEEILKSL (SEQ ID NO:30), VESYVPLFP (SEQ ID NO:31), FLLYQEPVLGPVR (SEQ ID NO:32), LNF (SEQ ID NO:33), and MHQPPQPLPPTVMFP (SEQ ID NO:34), an active analog thereof, and combinations thereof; wherein the active analog comprises a peptide having an amino acid sequence with at least about

15 percent proline and having at least about 70 percent ~~structural similarity~~ sequence identity to one or more constituent peptides of colostrinin, which are selected from the group of SEQ ID NO:1 through SEQ ID NO:34.

28. **(Previously Presented)** The method of claim 27 wherein the leukocyte regulator is selected from the group of MQPPPLP (SEQ ID NO:1), LQTPQPLLQVMMEPQGD (SEQ ID NO:2), DQPPDVEKPDQLQPFQVQS (SEQ ID NO:3), LFFFLPVVNVLP (SEQ ID NO:4), DLEMPVLPVEPFPPFV (SEQ ID NO:5), MPQNFYKLPQM (SEQ ID NO:6), VLEMKFPPPPQETVT (SEQ ID NO:7), LKPFPKLKVEVFPEP (SEQ ID NO:8), YPFTGPIPN (SEQ ID NO:18), SLPQNILPL (SEQ ID NO:19), TQTPVVVPPF (SEQ ID NO:20), HKEMPFPKYPVEPFOTESQ (SEQ ID NO:22), and combinations thereof.

29. **(Currently Amended)** A method for modulating leukocyte proliferation in a patient, the method comprising administering to the patient a leukocyte regulator selected from the group of colostrinin, a constituent peptide thereof, an analog thereof, and combinations thereof, under conditions effective to change the number of leukocytes; wherein the active analog comprises a peptide having an amino acid sequence with at least about 15 percent proline and having at least about 70 percent ~~structural similarity~~ sequence identity to one or more constituent peptides of colostrinin, which are selected from the group of SEQ ID NO:1 through SEQ ID NO:34.

30. **(Original)** The method of claim 29 wherein the patient is a human.

31. **(Previously Presented)** The method of claim 29 wherein the leukocytes are increased in number.

32. **(Previously Presented)** The method of claim 31 wherein the leukocytes are differentiated.

33. **(Previously Presented)** The method of claim 29 wherein the leukocyte regulator is a constituent peptide of colostrinin.

34. **(Currently Amended)** The method of claim 33 wherein the leukocyte regulator is selected from the group of MQPPPLP (SEQ ID NO:1), LQTPQPLLQVMMEPQGD (SEQ ID NO:2), DQPPDVEKPDQLQPFQVQS (SEQ ID NO:3), LFFFLPVVNVLP (SEQ ID NO:4), DLEMPVLPVEPFPPFV (SEQ ID NO:5), MPQNFYKLPQM (SEQ ID NO:6), VLEMKFPPPPQETVT (SEQ ID NO:7), LKPFPKLVVEVFPFP (SEQ ID NO:8), VVMEV (SEQ ID NO:9), SEQP (SEQ ID NO:10), DKE (SEQ ID NO:11), FPPPK (SEQ ID NO:12), DSQPPV (SEQ ID NO:13), DPPPPQS (SEQ ID NO:14), SEEMP (SEQ ID NO:15), KYKLQPE (SEQ ID NO:16), VLPPNVG (SEQ ID NO:17), VYPFTGPIPN (SEQ ID NO:18), SLPQNILPL (SEQ ID NO:19), TQTPVVVPPF (SEQ ID NO:20), LQPEIMGVPKVKETMVPK (SEQ ID NO:21), HKEMPFPKYPVEPFTESQ (SEQ ID NO:22), SLTLTDVEKLHLPLPLVQ (SEQ ID NO:23), SWMHQPP (SEQ ID NO:24), QPLPPTVMFP (SEQ ID NO:25), PQSVLS (SEQ ID NO:26), LSQPKVLPVPQKAVPQRDMPIQ (SEQ ID NO:27), AFLLYQE (SEQ ID NO:28), RGPFPILV (SEQ ID NO:29), ATFNRYQDDHGEEILKSL (SEQ ID NO:30), VESYVPLFP (SEQ ID NO:31), FLLYQEPVLGPVR (SEQ ID NO:32), LNF (SEQ ID NO:33), and MHQPPQPLPPTVMFP (SEQ ID NO:34), an active analog thereof, and combinations thereof; wherein the active analog comprises a peptide having an amino acid sequence with at least about 15 percent proline and having at least about 70 percent ~~structural similarity~~ sequence identity to one or more constituent peptides of colostrinin, which are selected from the group of SEQ ID NO:1 through SEQ ID NO:34.

35. **(Previously Presented)** The method of claim 34 wherein the leukocyte regulator is selected from the group of MQPPPLP (SEQ ID NO:1), LQTPQPLLQVMMEPQGD (SEQ ID NO:2), DQPPDVEKPDQLQPFQVQS (SEQ ID NO:3), LFFFLPVVNVLP (SEQ ID NO:4), DLEMPVLPVEPFPPFV (SEQ ID NO:5), MPQNFYKLPQM (SEQ ID NO:6),

VLEMKFPPPPQETVT (SEQ ID NO:7), LKPFPKLKVEVFPEP (SEQ ID NO:8), VYPFTGPIPN (SEQ ID NO:18), SLPQNILPL (SEQ ID NO:19), TQTPVVVPPF (SEQ ID NO:20), HKEMPFPKYPVEPFTESQ (SEQ ID NO:22), and combinations thereof.

36. **(Cancelled)**

37. **(Withdrawn-Currently Amended)** A method of inducing a cytokine in a cell, the method comprising contacting the cell with an immunological regulator under conditions effective to induce a cytokine, wherein the immunological regulator is a constituent peptide of colostrinin selected from the group consisting of LQTPQPLLQVMMEPQGD (SEQ ID NO:2), DQPPDVEKPDQLQPFQVQS (SEQ ID NO:3), LFFFLPVVNVLP (SEQ ID NO:4), DLEMPVLPVEFPFV (SEQ ID NO:5), MPQNFYKLPQM (SEQ ID NO:6), VLEMKFPPPPQETVT (SEQ ID NO:7), LKPFPKLKVEVFPPF (SEQ ID NO:8), VVMEV (SEQ ID NO:9), SEQP (SEQ ID NO:10), DKE (SEQ ID NO:11), FPPPK (SEQ ID NO:12), DSQPPV (SEQ ID NO:13), DPPPQS (SEQ ID NO:14), SEEMP (SEQ ID NO:15), KYKLQPE (SEQ ID NO:16), VLPPNVG (SEQ ID NO:17), VYPFTGPIPN (SEQ ID NO:18), SLPQNILPL (SEQ ID NO:19), TQTPVVVPPF (SEQ ID NO:20), LQPEIMGVPKVKETMVPK (SEQ ID NO:21), HKEMPFPKYPVEPFTESQ (SEQ ID NO:22), SLTLTDVEKLHLPLPLVQ (SEQ ID NO:23), SWMHQPP (SEQ ID NO:24), QLPPTVMFP (SEQ ID NO:25), PQSVLS (SEQ ID NO:26), LSQPKVLPVPQKAVPQRDMPIQ (SEQ ID NO:27), AFLLYQE (SEQ ID NO:28), RGPFPILV (SEQ ID NO:29), ATFNRYQDDHGEEILKSL (SEQ ID NO:30), FLLYQEPVLGPVR (SEQ ID NO:32), LNF (SEQ ID NO:33), and MHQPPQPLPPTVMFP (SEQ ID NO:34), an active analog thereof, and combinations thereof, wherein the active analog comprises a peptide having an amino acid sequence with at least about 15 percent proline and having at least about 70 percent ~~structural similarity~~ sequence identity to a one or more constituent peptides of colostrinin which are selected from the group of SEQ ID NO:2-30 and 32-34.

38. **(Withdrawn-Currently Amended)** A method for modulating an immune response in a cell, the method comprising contacting the cell with an immunological regulator under conditions effective to induce a cytokine, wherein the immunological regulator is a constituent peptide of colostrinin selected from the group consisting of LQTPQPLLQVMMEPQGD (SEQ ID NO:2), DQPPDVEKPDLPFQVQS (SEQ ID NO:3), LFFFLPVVNVLP (SEQ ID NO:4), DLEMPVLPVEPFPFV (SEQ ID NO:5), MPQNFYKLPQM (SEQ ID NO:6), VLEMKFPPPPQETVT (SEQ ID NO:7), LKPFPKLKVEVFPFP (SEQ ID NO:8), VVMEV (SEQ ID NO:9), SEQP (SEQ ID NO:10), DKE (SEQ ID NO:11), FPPPK (SEQ ID NO:12), DSQPPV (SEQ ID NO:13), DPPPPQS (SEQ ID NO:14), SEEMP (SEQ ID NO:15), KYKLQPE (SEQ ID NO:16), VLPPNVG (SEQ ID NO:17), VYPFTGPIPN (SEQ ID NO:18), SLPQNILPL (SEQ ID NO:19), TQTPVVVPPF (SEQ ID NO:20), LQPEIMGVVKVKETMVPK (SEQ ID NO:21), HKEMPFPKYPVEPFTESQ (SEQ ID NO:22), SLTLTDVEKLHLPLPLVQ (SEQ ID NO:23), SWMHQPP (SEQ ID NO:24), QPLPPTVMFP (SEQ ID NO:25), PQSVLS (SEQ ID NO:26), LSQPKVLPVPQKAVPQRDMPIQ (SEQ ID NO:27), AFLLYQE (SEQ ID NO:28), RGPFPILV (SEQ ID NO:29), ATFNRYQDDHGEEILKSL (SEQ ID NO:30), FLLYQEPVLGPVR (SEQ ID NO:32), LNF (SEQ ID NO:33), and MHQPPQPLPPTVMFP (SEQ ID NO:34), an active analog thereof, and combinations thereof, wherein the active analog comprises a peptide having an amino acid sequence with at least about 15 percent proline and having at least about 70 percent ~~structural-similarity~~ sequence identity to one or more constituent peptides of colostrinin which are selected from the group of SEQ ID NO:2-30 and 32-34.

39. **(Withdrawn-Currently Amended)** A method for modulating an immune response in a patient, the method comprising administering to the patient an immunological regulator under conditions effective to induce a cytokine, wherein the immunological regulator is a constituent peptide of colostrinin selected from the group consisting of LQTPQPLLQVMMEPQGD (SEQ ID NO:2), DQPPDVEKPDLPFQVQS (SEQ ID NO:3), LFFFLPVVNVLP (SEQ ID NO:4), DLEMPVLPVEPFPFV (SEQ ID NO:5), MPQNFYKLPQM (SEQ ID NO:6),



**Amendment and Response**

Page 10 of 18

Serial No.: 09/641,801

Confirmation No.: 5388

Filed: August 17, 2000

For: USE OF COLOSTRININ, CONSTITUENT PEPTIDES THEREOF, AND ANALOGS THEREOF FOR INDUCING CYTOKINES

---

VLEMKFPPPPQETVT (SEQ ID NO:7), LKPFPKLKVEVFPFP (SEQ ID NO:8), VVMEV (SEQ ID NO:9), SEQP (SEQ ID NO:10), DKE (SEQ ID NO:11), FPPPK (SEQ ID NO:12), DSQPPV (SEQ ID NO:13), DPPPPQS (SEQ ID NO:14), SEEMP (SEQ ID NO:15), KYKLQPE (SEQ ID NO:16), VLPPNVG (SEQ ID NO:17), VYPFTGPIPN (SEQ ID NO:18), SLPQNILPL (SEQ ID NO:19), TQTPVVVPPF (SEQ ID NO:20), LQPEIMGVPKVKETMVPK (SEQ ID NO:21), HKEMPFPKYPVEPFTESQ (SEQ ID NO:22), SLTLTDVEKLHLPLPLVQ (SEQ ID NO:23), SWMHQPP (SEQ ID NO:24), QPLPPTVMFP (SEQ ID NO:25), PQSVLS (SEQ ID NO:26), LSQPKVLPVPQKAVPQRDMPIQ (SEQ ID NO:27), AFLLYQE (SEQ ID NO:28), RGPFPILV (SEQ ID NO:29), ATFNRYQDDHGEEILKSL (SEQ ID NO:30), FLLYQEPVLGPVR (SEQ ID NO:32), LNF (SEQ ID NO:33), and MHQPPQPLPPTVMFP (SEQ ID NO:34), an active analog thereof, and combinations thereof, wherein the active analog comprises a peptide having an amino acid sequence with at least about 15 percent proline and having at least about 70 percent ~~structural similarity~~ sequence identity to one or more constituent peptides of colostrinin which are selected from the group of SEQ ID NO:2-30 and 32-34.